AD-A274 830

TION PAGE Form Approved OMB No. 0704-0188 Collection of information including suggestions for reducing this burgen to Davis Highway, Suite 1204. Artinator, VA 22202-4302 and to the Office of Ma for information Operations and Reported 10704-01881 Washington, DC, 20503 1. AGENCY USE ONLY (Leave blank) 2. REPORT DATE 3. REPORT TYPE AND DATES COVERED January 8, 1994 Technical Report 4. TITLE AND SUBTITLE 5. FUNDING NUMBERS Octahedral Metallomesogens:Liquid Crystallinity in Low Aspect Ratio **Materials** N00014-92-J-1416 6. AUTHOR(S) R&T Code 400X098YIP01 Hanxing Zheng and Timothy M. Swager 8. PERFORMING ORGANIZATION REPORT NUMBER 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Chemistry University of Pennsylvania Philadelphia, PA 19104-6323 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSORING/MONITORING AGENCY REPORT NUMBER Dr. H. Guard (Program Director)

Office of Naval Research 800 North Quincy Street Arlington, VA 22217-5000

1. SUPPLEMENTARY NOTES

J. Am. Chem. Soc. in press.

Technical Report No. 5

12a DISTRIBUTION/AVAILABILITY STATEMENT

Approved for public release; distribution unlimited

13. ABSTRACT (Maximum 200 words)

A new class of metallomesogens with octahedral structures are described. The observation of liquid crystallinity in these materials was unexpected as a result of their low aspect ratios. To better understand these materials, the dependence of the mesomorphism upon the number of sidechains (N) and metal center (Fe, Mn, Cr) was investigated. Complexes of Fe (N=6) appear to exhibit a highly ordered smectic phase. However, all of the complexes investigated with N=12, 15, and 18 display columnar mesomorphism. Columnar phases with hexagonal and rectangular symmetry were observed which display optical textures similar to known discotic phases. In addition it was found that the conformational preferences of the alkoxyphenyl groups attached to the 1,3-diketone ligands are coupled to the stability of different phases. Although these phases show textural and structural similarities with known discotic columnar phases, they are not miscible with discotic compounds having similar lattice parameters and regions of mesomorphism. It is proposed that the octahedral complexes exhibit clefts necessary to accommodate sidechains which can not be directed outside the columns core. As a result of the differences from regular discotics, these new columnar phases have been labeled as octahedral hexagonal Oh and octahedral rectangular Or.

14. SUBJECT TERMS			15, NUMBER OF PAGES
			18. PRICE CODE NA
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT
Unclassified	Unclassified	Unclassified	Unlimited

OFFICE OF NAVAL RESEARCH

Grant N00014-92-J-1416

R&T Code 400X098YIP01

Technical Report No. 5

Octahedral Metallomesogens: Liquid Crystallinity in Low Aspect Ratio Materials

by

Hanxing Zheng and Timothy M. Swager

J. Am. Chem. Soc. in press

Department of Chemistry University of Pennsylvania Philadelphia, PA 19104-6323

January 8, 1994

Reproduction in whole or in part is permitted for any purpose of the United States Government

This document has been approved for public release and sale; its distribution is unlimited

DIIO QUALITY INSPECTED B

Accesion For				
NTIS DTIC Unanno Justific	TAB ounced	d		
By Distrib	ution /			
Availability Codes				
Dist	Avail ar Spec			

Octahedral Metallomesogens: Liquid Crystallinity in Low Aspect Ratio Materials

Hanxing Zheng and Timothy M. Swager* Department of Chemistry University of Pennsylvania Philadelphia, PA 19104-6323

Abstract:

A new class of metallomesogens with octahedral structures 1, 2, and 3 are described. The observation of liquid crystallinity in these materials was unexpected as a result of their low aspect ratios. To better understand these materials, the dependence of the mesomorphism upon the number of sidechains (N) and metal center was investigated. Complexes of 1 (N=6) appear to exhibit a highly ordered smectic phase. However, all of the complexes investigated with N=12, 15, and 18 display columnar mesomorphism. Columnar phases with hexagonal and rectangular symmetry were observed which display optical textures similar to known discotic phases. In addition it was found that the conformational preferences of the alkoxyphenyl groups attached to the 1,3-diketone ligands are coupled to the stability of different phases. Although these phases show textural and structural similarities with known discotic columnar phases, they are not miscible with discotic compounds having similar lattice parameters and regions of mesomorphism. It is proposed that the octahedral complexes exhibit clefts necessary to accommodate sidechains which can not be directed outside the columns core. As a result of the differences from regular discotics, these new columnar phases have been labeled as octahedral hexagonal O_h and octahedral rectangular O_T.

$$R_{1}$$
 R_{2}
 R_{2}
 R_{3}
 R_{4}
 R_{5}
 R_{1}
 R_{2}
 R_{2}
 R_{3}
 R_{4}
 R_{5}
 R_{1}
 R_{2}
 R_{3}
 R_{1}
 R_{2}
 R_{3}
 R_{4}
 R_{5}
 R_{1}
 R_{2}
 R_{3}
 R_{1}
 R_{2}
 R_{3}
 R_{4}
 R_{5}
 R_{1}
 R_{1}
 R_{2}
 R_{3}
 R_{4}
 R_{5}
 R_{1}
 R_{1}
 R_{2}
 R_{3}
 R_{4}
 R_{5}
 R_{1}
 R_{2}
 R_{3}
 R_{4}
 R_{5}
 R_{1}
 R_{1}
 R_{2}
 R_{2}
 R_{3}
 R_{4}
 R_{5}
 R_{1}
 R_{2}
 R_{3}
 R_{4}
 R_{5}
 R_{1}
 R_{1}
 R_{2}
 R_{3}
 R_{4}
 R_{5}
 R_{1}
 R_{1}
 R_{2}
 R_{3}
 R_{4}
 R_{5}
 R_{5}
 R_{1}
 R_{1}
 R_{1}
 R_{2}
 R_{3}
 R_{4}
 R_{5}
 R_{5